

Reporting under the Habitat Regulations (as amended)¹

2019-2024

Conservation status assessment for the species:

S2030 - Risso's dolphin

(Grampus griseus)

United Kingdom



Scottish Government
Riaghaltas na h-Alba
gov.scot



Cyfoeth
Naturiol
Cymru
Natural
Resources
Wales



Department of
Agriculture, Environment
and Rural Affairs

An Roinn
Talmhaíochta, Comhshaoil
agus Gnóthaí Tuaithe

Department of
Fairmin, Environment
an' Kintra Matters

www.daera-ni.gov.uk



Northern Ireland Environment Agency
Gníomhaireacht Comhshaoil Thuaisceart Éireann
www.daera-ni.gov.uk | www.niea.gov.uk | www.norlin-airian-environment.gov.uk

¹ Habitat Regulations (as amended):

- The Conservation of Habitats and Species Regulations 2017 (as amended), Regulation 9A
- The Conservation of Offshore Marine Habitats and Species Regulations 2017 (as amended), Regulation 6A
- Report under The Conservation (Natural Habitats, &c.) Regulations 1994 (as amended), regulation 3ZA
- The Conservation (Natural Habitats, etc.) Regulations (Northern Ireland) 1995 (as amended), regulation 3ZA

For further information please contact:

Joint Nature Conservation Committee. Quay House, 2 East Station Road, Fletton Quays, Peterborough, PE2 8YY. <https://jncc.gov.uk>

This report was produced by JNCC in collaboration with the UK Country Nature Conservation Bodies (CNCBs) and country governments.

This document should be cited as:

JNCC, Department of Agriculture, Environment and Rural Affairs, Natural England, Natural Resources Wales & NatureScot. (2026). Conservation status assessment for the species: S2030 Risso's dolphin (*Grampus griseus*).

This resource and any accompanying material (e.g. maps, data, images) is published by JNCC under the [Open Government Licence](#) (OGLv3.0 for public sector information), unless otherwise stated. Note that some images (maps, tables) may not be copyright JNCC; please check sources for conditions of re-use.

The views and recommendations presented in this resource do not necessarily reflect the views and policies of JNCC.

Important note - Please read

- The information in this document represents the United Kingdom Reporting under the Habitat Regulations (as amended)¹, for the period 2019-2024.
- It is based on supporting information provided by Joint Nature Conservation Committee and UK Country Nature Conservation Bodies (CNCBs), which is documented separately.
- The Habitats Regulations reporting 2019-2024 Approach Document provides details on how this supporting information contributed to the UK Report and the fields that were completed for each parameter.
- Map showing the distribution and range of the species is included.
- Explanatory notes (where provided) are included at the end. These provide additional audit trail information to that included within the assessments. Further underpinning explanatory notes are available in the related country reports.
- Some of the reporting fields have been left blank because either: (i) there was insufficient information to complete the field; (ii) completion of the field was not obligatory; and/or (iii) the field was not relevant to this species (section 12 National Site Network coverage for Annex II species).

Further details on the approach to the Habitats Regulations Reporting 2019-2024 are available on the [JNCC website](#).

Assessment Summary: Risso's dolphin

Distribution and Range Map

Distribution and Range
Risso's dolphin




Figure 1: United Kingdom distribution and range map for S2030 - Risso's dolphin (*Grampus griseus*). The 50km grid square distribution map is based on available species records within the current reporting period.

Table 1: Table summarising the conservation status for S2030 - Risso's dolphin (*Grampus griseus*). Overall conservation status for species is based on assessments of range, population, habitat for the species, and future prospects.

Overall Conservation Status (see section 11)

Favourable (FV)

Breakdown of Overall Conservation Status

Range (see section 5)	Favourable (FV)
Population (see section 6)	Favourable (FV)
Habitat for the species (see section 7)	Unknown (XX)
Future prospects (see section 10)	Favourable (FV)

List of Sections

National Level	5
1. General information	5
2. Maps	5
3. Information related to Annex V Species	5
Biogeographical Level	7
4. Biogeographical and marine regions	7
5. Range	7
6. Population	9
7. Habitat for the species	12
8. Main pressures	13
9. Conservation measures	14
10. Future prospects	15
11. Conclusions	16
12. UK National Site Network (pSCIs, SCIs, SACs) coverage for Annex II species	18
13. Complementary information	18
14. References	20
Biogeographical and marine regions	20
Main pressures	24
15. Explanatory Notes	25

National Level

1. General information

1.1 Country	United Kingdom
1.2 Species code	S2030
1.3 Species scientific name	<i>Grampus griseus</i>
1.4 Alternative species scientific name	
1.5 Common name	Risso's dolphin
Annex(es)	IV

2. Maps

2.1 Sensitive species	No
2.2 Year or period	2019-2022
2.3 Distribution map	Yes
2.4 Distribution map; Method used	Complete survey or a statistically robust estimate

2.5 Additional information

The distribution map is based on verified sightings data of Risso's dolphin between 2019 and 2024. The sightings were collated from SCANS IV, Pelagis French surveys, NBN Atlas, European Seabirds at Sea, the Joint Cetacean Data Programme, POSEIDON project, University of Aberdeen, The Crown Estate Marine Data Exchange, Whale and Dolphin Conservation, Hebridean Whale and Dolphin Trust, ORCA, Sea Watch Foundation, Marine Discovery Penzance, Sussex Dolphin Project, Cornwall Seal Group Research Trust, Cardigan Bay Marine Wildlife Centre, Natural England, Sea Trust and The Royal Society for the Protection of Birds (RSPB).

3. Information related to Annex V Species

3.1 Is the species taken in the wild / exploited?

3.2 What measures have been taken?

a) Regulations regarding access to property

b) Temporary or local prohibition on the taking of specimens in the wild and exploitation

c) Regulation of the periods and/or methods of taking specimens

d) Application of hunting and fishing rules which take account of the conservation of such populations

e) Establishment of a system of licences for taking specimens or of quotas

f) Regulation of the purchase, sale, offering for sale, keeping for sale, or transport for sale of specimens

g) Breeding in captivity of animal species as well as artificial propagation of plant species

Other measures

Other measures description

3.3: Hunting bag or quantity taken in the wild for Mammals and Acipenseridae (Fish)

a) Unit

Table 2: Quantity taken from the wild during the reporting period (see 3.3a for units). For species with defined hunting seasons, Season 1 refers to 2018/2019 (autumn 2018 to spring 2019), and Season 6 to 2023/2024. For species without hunting seasons, data are reported by calendar year: Year 1 is 2019, and Year 6 is 2024.

	Season/ year 1	Season/ year 2	Season/ year 3	Season/ year 4	Season/ year 5	Season/ year 6
b) Minimum	-	-	-	-	-	-
c) Maximum	-	-	-	-	-	-
d) Unknown	-	-	-	-	-	-

3.4: Hunting bag or quantity taken in the wild; Method used

3.5: Additional information

No additional information

Biogeographical Level

4. Biogeographical and marine regions

4.1 Biogeographical or marine region where the species occurs MATL

4.2 Sources of information

See section 14 References

5. Range

5.1 Surface area (km²) 565,632

5.2 Short-term trend; Period 2013-2024

5.3 Short-term trend; Direction Increasing

5.4 Short-term trend; Magnitude

a) Estimated minimum

b) Estimated maximum

c) Pre-defined range

d) Unknown

e) Type of estimate

f) Rate of decrease

5.5 Short-term trend; Method used Complete survey or a statistically robust estimate used

5.6 Long-term trend; Period

5.7 Long-term trend; Direction

**5.8 Long-term trend;
Magnitude**

a) Minimum

b) Maximum

c) Rate of decrease

**5.9 Long-term trend; Method
used**

5.10 Favourable Reference Range (FRR)

a) Area (km²) 531,659

b) Pre-defined increment

c) Unknown No

d) Method used Model-based approach

e) Quality of information high

5.11 Change and reason for change in surface area of range

a) Change Yes

b) Genuine change No

**c) Improved knowledge or
more accurate data** Yes

d) Different method No

e) No information No

f) Other reason No

g) Main reason Improved knowledge/more accurate data

5.12 Additional information

The distribution is based on verified sightings data of Risso's dolphin between 2019 and 2024. The sightings were collated from SCANS IV, Pelagis French surveys, NBN Atlas, European Seabirds at Sea, the Joint Cetacean Data Programme, POSEIDON project, University of Aberdeen, The Crown Estate Marine Data Exchange, Whale and Dolphin Conservation, Hebridean Whale and Dolphin Trust, ORCA, Sea Watch Foundation, Marine Discovery Penzance, Sussex Dolphin Project, Cornwall Seal Group Research

Trust, Cardigan Bay Marine Wildlife Centre, Natural England, Sea Trust and The Royal Society for the Protection of Birds (RSPB).

The FRR was based on an analysis of effort related survey data spanning 1994-2010 compiled for the Joint Cetacean Protocol (JCP) undertaken by Paxton et al. (2016). The estimated range was based on a modelled prediction of Risso's dolphin distribution during August 2010 and adapted based on additional sightings data and expert knowledge (see Paxton et al., 2016 for further detail).

Based on recent modelling of cetacean species (such as Waggitt et al., 2019) and the distribution of observations from citizen science and systematic surveys, Risso's dolphin occur in deeper waters of the English Channel, though in lower densities than other regions. Therefore, the range has been updated to include areas where Risso's have been recorded but were not covered by the 2013 modelled range. This change reflects improvement in our knowledge of this species, rather than a genuine change.

Since the 2019 Habitats Directive Article 17 assessments, the FRR has changed due to the removal of the EEZ extension into offshore waters west of Scotland. This area has been removed due to lack of data for all species, and subsequent impact on confidence in assessments. This does not represent genuine change in FRR.

6. Population

6.1 Year or period 2022

6.2 Population size (in reporting unit)

a) Unit number of individuals

b) Minimum 3,389

c) Maximum 24,307

d) Best single value 9,076

6.3 Type of estimate 95% confidence interval

6.4 Quality of extrapolation to reporting unit moderate

6.5 Additional population size (using population unit other than reporting unit)

a) Unit

b) Minimum

c) Maximum

d) Best single value

e) Type of estimate

6.6 Population size; Method used Complete survey or a statistically robust estimate used

6.7 Short-term trend; Period 2016-2022

6.8 Short-term trend; Direction Stable

6.9 Short-term trend; Magnitude

a) Estimated minimum

b) Estimated maximum

c) Pre-defined range

d) Unknown

e) Type of estimate

f) Rate of decrease

6.10 Short-term trend; Method used Complete survey or a statistically robust estimate used

6.11 Long-term trend; Period

6.12 Long-term trend; Direction Unknown

6.13 Long-term trend; Magnitude

a) Minimum

b) Maximum

c) Confidence interval

d) Rate of decrease

6.14 Long-term trend; Method used Insufficient or no data available

6.15 Favourable Reference Population (FRP)

ai) Population size	8,917
aii) Unit	number of individuals
b) Pre-defined increment	
c) Unknown	No
d) Method used	Model-based approach
e) Quality of information	moderate

6.16 Change and reason for change in population size

a) Change	No
b) Genuine change	
c) Improved knowledge or more accurate data	
d) Different method	
e) No information	
f) Other reason	
g) Main reason	

6.17 Additional information

There are only two abundance estimates available for Risso's dolphin within the UK EEZ from SCANS III (Hammond et al. 2021) and SCANS IV (Gilles, et al., 2023) which limits capacity to confidently identify trends. The current evidence suggests the population may be stable, however due to the high confidence intervals surrounding abundance estimates for Risso's dolphin, there is limited statistical power to detect anything beyond a dramatic change in abundance.

In terms of the wider context for Risso's dolphin population in the NE Atlantic; ObSERVE in Irish waters found an increase in abundance between 2015 and 2022, especially in winter months (Giralt Paradell, et al. 2024). Risso's dolphin are a wide ranging species, found between Norwegian waters to the west coast of Africa. Findings from NASS 2024 survey in the NAMMCO region will provide more context for the northern areas of their range, once published.

The FRV for population was calculated based on estimates from SCANS III (Hammond et al., 2021), supplemented with density estimates from neighbouring regions to fill data gaps within the UK EEZ and limit extrapolation where possible; ObSERVE in Irish

waters (Rogan, e al., 2018), NASS 2015 (Pike, et al., 2019a) and NILS 2015 (Leonard and Øien, 2020a) surveys in the NAMMCO region.

Since the 2019 Habitats Directive Article 17 assessments, the FRV has changed due to the removal of the EEZ extension into offshore waters west of Scotland. This area has been removed due to lack of data for all species, and subsequent impact on confidence in assessments. This does not represent genuine change in FRV.

6.18 Age structure, mortality and reproduction deviation Unknown

7. Habitat for the species

7.1 Sufficiency of area and quality of occupied habitat (for long-term survival)

a) Is area of occupied habitat sufficient? Unknown

b) Is quality of occupied habitat sufficient? Unknown

c) If No or Unknown, is there a sufficiently large area of unoccupied habitat of suitable quality? Unknown

7.2 Sufficiency of area and quality of occupied habitat; Method used

a) Sufficiency of area of occupied habitat; Method used Based mainly on expert opinion with very limited data

b) Sufficiency of quality of occupied habitat; Method used Based mainly on expert opinion with very limited data

7.3 Short-term trend; Period

7.4 Short-term trend; Direction Unknown

7.5 Short-term trend; Method used Based mainly on expert opinion with very limited data

7.6 Long-term trend; Period

7.7 Long-term trend; Direction Unknown

7.8 Long-term trend; Method used

Based mainly on expert opinion with very limited data

7.9 Additional information

Direct evidence of cetacean habitat quality is limited as presently, a comprehensive understanding of the key elements important to the species is undetermined. In some cases, conclusions for species range and population could be indicative of habitat quality by proxy, however confidence in assessment outputs would be low.

The population of Risso’s dolphin using the UK EEZ is stable since 2016 and the range has also remained stable.

8. Main pressures

8.1 Characterisation of pressures

Table 3: Pressures affecting the species, including timing and importance/impact ranking. Pressures are defined as factors acting currently and/or during the reporting period (2019–2024). Rankings are: High (direct/immediate influence and/or large spatial extent) and Medium (moderate direct/immediate influence, mainly indirect and/or regional extent).

Pressure	Timing	Ranking
Wind, wave and tidal power (including infrastructure)	Ongoing and likely to be in the future	Medium (M)
Sports, tourism and leisure activities	Ongoing and likely to be in the future	Medium (M)
Residential, commercial and industrial activities and structures generating noise, light, heat or other forms of pollution	Ongoing and likely to be in the future	Medium (M)
Bycatch and incidental killing (due to fishing and hunting activities)	Ongoing and likely to be in the future	Medium (M)
Mixed source marine water pollution (marine and coastal)	Ongoing and likely to be in the future	Medium (M)
Decline or extinction of related species (e.g. food source / prey, predator / parasite, symbiote, etc.) due to climate change	Ongoing and likely to be in the future	High (H)
Geotechnical surveying	Ongoing and likely to be in the future	Medium (M)

8.2 Sources of information

See section 14 References

8.3 Additional information

PD01: Greater concern in West Scotland and Wales.

PF05: Assessed as a 'Medium' pressure but current reporting may be underestimating the extent of the pressure on marine mammal populations around the UK. A greater pressure in coastal areas.

PC07: Regional pressure in the North Sea and the Irish Sea.

9. Conservation measures

9.1: Status of measures

a) Are measures needed?

Yes

b) Indicate the status of measures

Measures identified and taken

9.2 Main purpose of the measures taken

Maintain the current range, population and/or habitat for the species

9.3 Location of the measures taken

Both inside and outside National Site Network

9.4 Response to measures

Medium-term results (within the next two reporting periods, 2025–2036)

9.5 List of main conservation measures

Table 4: Key conservation measures addressing current pressures and/or anticipated threats during the next two reporting periods (2025–2036). Measures are ranked by importance/impact: High (direct/immediate influence and/or large spatial extent) and Medium (moderate direct/immediate influence, mainly indirect and/or regional extent).

Conservation measure	Ranking
Adapt/manage exploitation of energy resources	High (H)
Control/eradication of illegal killing, fishing and harvesting of wild plants, fungi and animals	High (H)
Reduce bycatch and incidental killing of non-target species	High (H)
Reduce impact of military installations and activities	High (H)

Reduce impact of mixed source pollution	High (H)
Adapt/manage renewable energy installation, facilities and operation (excl. hydropower and abstraction activities)	High (H)
Management of professional/commercial fishing, shellfish and seaweed harvesting (incl. restoration of habitats)	High (H)

9.6 Additional information

This species is not an Annex II species and therefore the designation of SACs is not required, as stipulated in the Habitats Regulations. However, as a European Protected Species, protection is provided throughout UK waters and it is an offence to kill, injure or disturb. The UK remains committed to the conservation of marine mammals in UK waters and the implementation of measures to mitigate the impact of pressures and conservation measures have been undertaken in the UK and adjacent waters as part of the requirements of the Habitats Regulations. Such measures include monitoring bycatch, monitoring strandings data to monitor current and identify emerging pressures, application of appropriate management measures, and noise monitoring and mitigation with regards to offshore industry. This is reflected in the list of conservation measures under field 9.5. The UK also supports a range of international agreements and conventions on the conservation of marine mammals and the marine environment. For example: The Convention on Migratory Species; the Convention for the Protection of the Marine Environment of the North-East Atlantic (OSPAR). A Marine Protected Area (MPA) has been designated with Risso's dolphin as a qualifying feature (grade A-C): (555703753) North-east Lewis MPA. A UK Cetacean Conservation Strategy is currently in development, due for publication shortly. The strategy is intended to support decision making and identify actions necessary to maintain or improve the conservation status of cetaceans in UK waters. Defra and devolved administrations fund national strandings schemes for cetaceans which aim to: collate, analyse and report data for all cetacean strandings around the coast of the UK; determine the causes of death (both natural and anthropogenic) in stranded cetaceans, including bycatch and physical trauma and; undertake surveillance on the incidence of disease in stranded cetaceans in order to identify any substantial new threats to their conservation status. The UK has several voluntary wildlife watching guidelines/codes of conduct which are publicly available however, while these are endorsed by the UK government and devolved administrations, there is no mandate to adopt by operators or individuals.

10. Future prospects

10.1a Future trends of parameters

ai) Range	Overall stable
bi) Population	Overall stable
ci) Habitat for the species	Unknown

10.1b Future prospects of parameters

aii) Range	Good
bii) Population	Good
cii) Habitat for the species	Unknown

10.2 Additional information

No additional information

11. Conclusions

11.1 Range	Favourable (FV)
11.2 Population	Favourable (FV)
11.3 Habitat for the species	Unknown (XX)
11.4 Future prospects	Favourable (FV)
11.5 Overall assessment of Conservation Status	Favourable (FV)
11.6 Overall trend in Conservation Status	Stable

11.7 Change and reason for change in conservation status

a) Change	Yes
b) Genuine change	No
c) Improved knowledge or more accurate data	Yes
d) Different method	No
e) No information	No
f) Other reason	No

g) Main reason	Improved knowledge/more accurate data
-----------------------	---------------------------------------

11.7 Change and reason for change in conservation status trend

a) Change	Yes
------------------	-----

b) Genuine change	No
--------------------------	----

c) Improved knowledge or more accurate data	Yes
----------------------------------------------------	-----

d) Different method	No
----------------------------	----

e) No information	No
--------------------------	----

f) Other reason	No
------------------------	----

g) Main reason	Improved knowledge/more accurate data
-----------------------	---------------------------------------

11.8 Additional information

Conclusion on Range reached because: (i) the short-term trend direction in Range surface area is increasing and (ii) the current Range surface area is approximately equivalent to the Favourable Reference Range.

Conclusion on Population reached because: (i) the short-term trend direction in Population size is stable; and (ii) the best estimate for population size is greater than the Favourable Reference Population.

Conclusion on Habitat for the species reached because: (i) it is unknown whether the area of habitat is sufficiently large; (ii) it is unknown if habitat quality is sufficient for the long-term survival of the species; and (iii) the short-term trend in area and quality of habitat is unknown.

Conclusion on Future prospects reached because: (i) the Future prospects for Range are Good; (ii) the Future prospects for Population are Good; and (iii) the Future prospects for Habitat for the species are Unknown.

Overall assessment of Conservation Status is Favourable because all conclusions, except for Habitat, are Favourable.

Overall trend in Conservation Status is based on the combination of the short-term trends for Range - increasing, Population - stable, and Habitat for the species - unknown.

12. UK National Site Network (pSCIs, SCIs, SACs) coverage for Annex II species

12.1 Population size inside the pSCIs, SCIs and SACs network

a) Unit

b) Minimum

c) Maximum

d) Best single value

12.2 Type of estimate

12.3 Population size inside the network; Method used

12.4 Short-term trend of population size within the network; Direction

12.5 Short-term trend of population size within the network; Method used

12.6 Short-term trend of habitat for the species inside the pSCIs, SCIs and SACs network; Direction

12.7 Short-term trend of habitat for the species inside the pSCIs, SCIs and SACs network; Method used

12.8 Additional information

No additional information

13. Complementary information

13.1 Justification of percentage thresholds for trends

No justification information

13.2 Trans-boundary assessment

No trans-boundary assessment information

13.2 Other relevant information

No other relevant information

14. References

Biogeographical and marine regions

4.2 Sources of information

Paxton, C.G.M., Scott-Hayward, L., Mackenzie, M., Rexstad, E. & Thomas, L. (2016) Revised Phase III Data Analysis of Joint Cetacean Protocol Data Resource, JNCC Report No. 517, JNCC, Peterborough, ISSN 0963-8091. <https://hub.jncc.gov.uk/assets/01adfabd-e75f-48ba-9643-2d594983201e>

Waggitt, J.J., Evans, P.G.H., Andrade, J., Banks, A.N., Boisseau, O., Bolton, M., Bradbury, G., Brereton, T., Camphuysen, C.J., Durinck, J., Felce, T., Fijn, R.C., Garcia-Baron, I., Garthe, S., Geelhoed, S.C.V., Gilles, A., Goodall, M., Haelters, J., Hamilton, S., Hartny-Mills, L., Hodgins, N., James, K., Jessopp, M., Kavanagh, A.S., Leopold, M., Lohrengel, K., Louzao, M., Markones, N., Martínez-Cedeira, J., Ó Cadhla, O., Perry, S.L., Pierce, G.J., Ridoux, V., Robinson, K.P., Santos, M.B., Saavedra, C., Skov, H., Stienen, E.W.M., Sveegaard, S., Thompson, P., Vanermen, N., Wall, D., Webb, A., Wilson, J., Wanless, S., Hiddink, J.G., 2020. Distribution maps of cetacean and seabird populations in the North-East Atlantic. *Journal of Applied Ecology* 57, 253–269. <https://doi.org/10.1111/1365-2664.13525>

Hammond, PS, Lacey, C, Gilles, A, Viquerat, S, Börjesson, P, Herr, H, Macleod, K, Ridoux, V, Santos, MB, Scheidat, M, Teilmann, J, Vingada, J & Øien, N (2021). Estimates of cetacean abundance in European Atlantic waters in summer 2016 from the SCANS-III aerial and shipboard surveys. SCANS-III project report 1, 39 pp. https://scans3.wp.st-andrews.ac.uk/files/2021/06/SCANS-III_design-based_estimates_final_report_revised_June_2021.pdf

Gilles, A, Authier, M, Ramirez-Martinez, NC, Araújo, H, Blanchard, A, Carlström, J, Eira, C, Dorémus, G, Fernández-Maldonado, C, Geelhoed, SCV, Kyhn, L, Laran, S, Nachtsheim, D, Panigada, S, Pigeault, R, Sequeira, M, Sveegaard, S, Taylor, NL, Owen, K, Saavedra, C, Vázquez-Bonales, JA, Unger, B, Hammond, PS (2023). Estimates of cetacean abundance in European Atlantic waters in summer 2022 from the SCANS-IV aerial and shipboard surveys. Final report published 29 September 2023. 64 pp. <https://www.tiho-hannover.de/itaw/scans-iv-survey>

Giralt Paradell, O., Cañadas, A., Bennison, A., Todd, N., Jessopp, M., Rogan, E. (2024). Aerial

surveys of cetaceans and seabirds in Irish waters: Occurrence, distribution and abundance in

2021-2023. Department of the Environment, Climate & Communications and
Department of

Housing, Local Government & Heritage, Ireland. 260pp [https://www.gov.ie/pdf/?file=
https://assets.gov.ie/308027/e03a534c-0fa5-4a22-8bad-5f002ae94857.pdf](https://www.gov.ie/pdf/?file=https://assets.gov.ie/308027/e03a534c-0fa5-4a22-8bad-5f002ae94857.pdf)

Bearzi, G., Reeves, R.R., Remonato, E., Pierantonio, N. and Airoidi, S., 2011. Risso's dolphin *Grampus griseus* in the Mediterranean Sea. *Mammalian Biology*, 76(4), pp.385-400.

Blanco, C., Raduán, M.Á. and Raga, J.A., 2006. Diet of Risso's dolphin (*Grampus griseus*) in the western Mediterranean Sea. *Scientia Marina*, 70(3), pp.407-411.

David, L., Arcangeli, A., Donovan, G., Holcer, D., Lanfredi, C., Airoidi, S., Dhermain, F., Fossi, C., Labach, H., Natoli, A., Pavan, G. Rosso, M. 2021. Progress report regarding Risso dolphin conservation management plan (CMP) in ACCOBAMS area. Available at: https://accobams.org/wp-content/uploads/2021/09/SC14.Doc16_Progress-Report-CMP-Risso.pdf. [Accessed 07 Nov 2024]

Fernandez-Betelu, O., Graham, I.M., Malcher, F., Webster, E., Cheong, S.H., Wang, L., Iorio-Merlo, V., Robinson, S. and Thompson, P.M., 2024. Characterising underwater noise and changes in harbour porpoise behaviour during the decommissioning of an oil and gas platform. *Marine Pollution Bulletin*, 200, p.116083.

Hermannsen, L., Beedholm, K., Tougaard, J. and Madsen, P.T., 2014. High frequency components of ship noise in shallow water with a discussion of implications for harbor porpoises (*Phocoena phocoena*). *The Journal of the Acoustical Society of America*, 136(4), pp.1640-1653.

Jepson, P.D., Deaville, R., Barber, J.L., Aguilar, À., Borrell, A., Murphy, S., Barry, J., Brownlow, A., Barnett, J., Berrow, S. and Cunningham, A.A., 2016. PCB pollution continues to impact populations of orcas and other dolphins in European waters. *Scientific reports*, 6(1), pp.1-17.

JNCC. 2010a. The protection of marine European Protected Species from deliberate injury, killing and disturbance. Guidance for the marine area in England and Wales and the UK offshore marine area. Available on request from JNCC.

JNCC. 2010b. Statutory nature conservation agency protocol for minimising the risk of injury to marine mammals from Piling noise. 2010. JNCC Peterborough. United Kingdom. Available at: <https://data.jncc.gov.uk/data/31662b6a-19ed-4918-9fab-8fbcff752046/JNCC-CNCB-Piling-protocol-August2010-Web.pdf> [Accessed 06 Nov 2024]

JNCC. 2010c. JNCC guidelines for minimising the risk of injury to marine mammals from using explosives. August 2010. Available at: <https://data.jncc.gov.uk/data/24cc180d->

[4030-49dd-8977-a04ebe0d7aca/JNCC-Guidelines-Explosives-Guidelines-201008-Web.pdf](https://data.jncc.gov.uk/data/e2a46de5-43d4-43f0-b296-c62134397ce4/jncc-guidelines-seismicsurvey-aug2017-web.pdf) [Accessed 06 Nov 2024]

JNCC. 2017. JNCC guidelines for minimising the risk of injury to marine mammals from geophysical surveys. Available at: <https://data.jncc.gov.uk/data/e2a46de5-43d4-43f0-b296-c62134397ce4/jncc-guidelines-seismicsurvey-aug2017-web.pdf> [Accessed 06 Nov 2024]

JNCC. 2023. JNCC guidance for the use of Passive Acoustic Monitoring in UK waters for minimising the risk of injury to marine mammals from offshore activities. JNCC, Peterborough. Available at: <https://hub.jncc.gov.uk/assets/fb7d345b-ec24-4c60-aba2-894e50375e33> [Accessed 06 Nov 2024]

Kim, G.B., Tanabe, S., Iwakiri, R., Tatsukawa, R., Amano, M., Miyazaki, N. and Tanaka, H., 1996. Accumulation of butyltin compounds in Risso's dolphin (*Grampus griseus*) from the Pacific coast of Japan: comparison with organochlorine residue pattern. *Environmental science & technology*, 30(8), pp.2620-2625.

Malinka, C.E., Gillespie, D.M., Macaulay, J.D., Joy, R. and Sparling, C.E., 2018. First in situ passive acoustic monitoring for marine mammals during operation of a tidal turbine in Ramsey Sound, Wales. *Marine Ecology Progress Series*, 590, pp.247-266.

Marine Scotland. 2014. The protection of Marine European Protected Species from injury and disturbance. Guidance for Scottish Inshore Waters.

Marsili, L. and Focardi, S., 1997. Chlorinated hydrocarbon (HCB, DDTs and PCBs) levels in cetaceans stranded along the Italian coasts: an overview. *Environmental Monitoring and Assessment*, 45, pp.129-180.

Megson, D., Brown, T., Jones, G.R., Robson, M., Johnson, G.W., Tiktak, G.P., Sandau, C.D. and Reiner, E.J., 2022. Polychlorinated biphenyl (PCB) concentrations and profiles in marine mammals from the North Atlantic Ocean. *Chemosphere*, 288, p.132639.

Middel, H. and Verones, F., 2017. Making marine noise pollution impacts heard: the case of cetaceans in the North sea within life cycle impact assessment. *Sustainability*, 9(7), p.1138.

Minoia, L., Consales, G., Mazzariol, S., Mancusi, C., Terracciano, G., Ceciari, I., Capanni, F., Neri, A., D'Agostino, A. and Marsili, L., 2023. Preliminary assessment of persistent organic pollutants (POPs) in tissues of Risso's dolphin (*Grampus griseus*) specimens stranded along the Italian coasts. *Marine Pollution Bulletin*, 186, p.114470.

Pierce, G.J., Brownlow, A., Evans, P.G., IJsseldijk, L., Kamińska, K., Kessler, L., Murphy, S., Pinn, E., Ridoux, V., Simmonds, M.P. and Spitz, J., 2022. Report of the ASCOBANS Resource Depletion Working Group. Available at: <https://www.ascobans.org/sites/>

default/files/document/ascobans_ac27_doc2.2_report-resource-depletion-wg.pdf

[Accessed 07 Nov 2024]

Plint, T., ten Doeschate, M.T., Brownlow, A.C., Davison, N.J., Hantke, G., Kitchener, A.C., Longstaffe, F.J., McGill, R.A., Simon-Nutbrown, C. and Magill, C.R., 2023. Stable isotope ecology and interspecific dietary overlap among dolphins in the Northeast Atlantic. *Frontiers in Marine Science*, 10, p.1111295.

Scottish Marine Animal Stranding Scheme. 2022. Scottish Marine Animal Stranding Scheme (SMASS) Annual Report 2021. Available at: <https://strandings.org/wp-content/uploads/2024/06/SMASS-Annual-Report-2021-final.pdf> [Accessed 06 Nov 2024]

Scottish Marine Animal Stranding Scheme. 2023. Scottish Marine Animal Stranding Scheme (SMASS) Annual Report 2022. Available: <https://strandings.org/wp-content/uploads/2024/06/SMASS-Annual-Report-2022-v1.2.pdf> [Accessed 07 Nov 2024]

Stone, C., Hall, K., Mendes, S. and Tasker, M. 2017. The effects of seismic operations in UK waters: analysis of Marine Mammal Observer data. *J. Cetacean Res. Manage.*, 16, pp.71-85.

Stone, C.J. 2015. Implementation of and considerations for revisions to the JNCC guidelines for seismic surveys. JNCC Report No. 463b. Available at: <https://data.jncc.gov.uk/data/f7990481-7a99-414c-be04-b972da10c1b7/JNCC-Report-463b-FINAL-WEB.pdf> [Accessed 06 Nov 2024]

Storelli, M.M., Zizzo, N. and Marcotrigiano, G.O., 1999. Heavy metals and methylmercury in tissues of Risso's dolphin (*Grampus griseus*) and Cuvier's beaked whale (*Ziphius cavirostris*) stranded in Italy (South Adriatic sea).

Thomsen, F., Ram, M., Chreptowicz, M., Nocoń, M. and Balicka, I., 2023. Noise modelling and environmental risk assessment of a geophysical survey and its impact on herring and minke whales in Irish coastal waters. Marine Institute.

Visser, F., Hartman, K.L., Rood, E.J., Hendriks, A.J., Zult, D.B., Wolff, W.J., Huisman, J. and Pierce, G.J., 2011. Risso's dolphins alter daily resting pattern in response to whale watching at the Azores. *Marine Mammal Science*, 27(2), pp.366-381.

JNCC. 2025. JNCC guidelines for minimising the risk of injury to marine mammals from unexploded ordnance (UXO) clearance in the marine environment. JNCC, Aberdeen.

JNCC, Natural England and Cefas. 2025. JNCC, Natural England and Cefas position on the use of quieter piling methods and noise abatement systems when installing offshore wind turbine foundations. JNCC, Aberdeen.

Evans, P.G.H. and Waggitt, J.J. 2023. Modelled Distribution and Abundance of Cetaceans and Seabirds in Wales and Surrounding Waters. NRW Evidence Report, Report No: 646,

354 pp. Natural Resources Wales, Bangor.

Main pressures

8.2 Sources of information

No sources of information

15. Explanatory Notes

Field label	Note
8.1: Characterisation of pressures	<p>PF12 Industrial or commercial activities and structures generating noise, light, heat or other forms of pollution. Cetaceans rely on echolocation for navigation, foraging and communication, making them sensitive to noise pollution in the marine environment (Middel and Verones, 2017; David et al., 2021). Noise pollution from shipping vessels is of concern to the species, as a diverse range of vessels produce substantial noise across a broad spectrum of frequencies (0.025-160kHz), including high frequencies where toothed whale hearing is most sensitive. Vessel noise should be considered over a broad frequency range, and not just low frequency bands, when assessing noise on effects on toothed whales (Hermanssen et al, 2014). Although alone this pressure is not graded Medium or High, the cumulative impact of this and other sources of noise may be greater when combined, which is accounted for here.</p>
8.1: Characterisation of pressures	<p>PF05 Sports, tourism and leisure activities. The impact of this pressure is indirect, and recovery from disturbance is unknown. Exposure is limited both spatially and temporally, although it may be regionally significant when occurring in areas known to be favoured by the species. Boat presence is associated with a short-term reduction in resting behaviour in Risso's dolphins. Disturbance from tourism activities impacts distribution and communication between the species and increases with frequency and number of vessels present (Visser et al., 2011; David et al., 2021). This pressure is mitigated to some degree by codes of conduct regarding interactions.</p>
8.1: Characterisation of pressures	<p>PD01 Wind, wave and tidal power, including infrastructure. There is limited evidence for the current effects of this pressure on Risso's dolphin. There are considerable legal and societal obligations to meet clean energy requirements which will result in an increase in the development of the renewable energy industry such as tidal and wave power,</p>

	with the potential to increase risk of collision (Malinka et al., 2018) and displacement from key habitat. This is regionally significant for Risso's dolphin, resulting in a Medium grading.
8.1: Characterisation of pressures	PJ12 Decline or extinction of related species (e.g. food source / prey, predator / parasite, symbiot, etc.) due to climate change. There is no current evidence for the effects of climate change on related species regarding Risso's dolphin. However, the species has a specific diet with a preference for cephalopods (Blanco et al., 2006; Bearzi et al., 2011; Pierce et al., 2022; Plint et al., 2023) which may reduce capacity to adapt to changes in prey availability as a result of changes in climate. Furthermore, habitat degradation from anthropogenic activities may exacerbate the issue around depletion of prey (David et al., 2021)
8.1: Characterisation of pressures	PK02 Mixed source marine water pollution (marine and coastal). This pressure has an indirect effect on mortality, mediated through the diet (bioaccumulation), causing reduced resilience to disease and lower fecundity through increased foetal mortality. The influence is long-term and intergenerational, with the pressure ubiquitous across the species range (Jepson et al, 2016). Evidence of pollutant burdens in Risso's dolphins exists for various populations globally (Storelli et al., 1999, Kim et al., 1996, Marsili & Focardi, 1997) and recent analysis in the UK indicates the issue is persisting (Jepson et al., 2016; Megson et al., 2022; Minoia et al., 2023). 155 different PCBs in Risso's dolphin were found during recent analysis of samples from the North Atlantic, the largest number found, and concentrations generally exceeded toxic thresholds (Megson et al., 2022). Other analyses have also found evidence of HCBs and DDTs in tissues of Risso's dolphin (Minoia et al., 2023). Evidence as to the impacts on this species are limited.
8.1: Characterisation of pressures	PG13 Bycatch and incidental killing (due to fishing and hunting activities). Cause of death in three stranded animals was attributed to bycatch or entanglement between 2018 and 2022 (30% of animals examined at post mortem;

	<p>Scottish Marine Animal Stranding Scheme 2022; 2023). This pressure was ranked Medium as it is recognised as potentially having an important direct and immediate influence on individuals, but with relatively low exposure.</p>
<p>8.1: Characterisation of pressures</p>	<p>PC07 Geotechnical surveying. JNCC advice on geophysical surveying considers the risk of activities to all marine mammals in UK waters (Stone, 2015; JNCC, 2017, 2010b, 2010c) and strong avoidance reactions and behavioural changes have been well documented for other odontocetes (Stone, 2015; Stone et al., 2017; Fernandez-Betelu et al., 2021; Thomsen et al., 2023). Furthermore, reviews on the threats faced by Risso's dolphins in other regions highlight the risk of impulsive noise from seismic surveys (David et al., 2021) and thus, it seems likely that the pressure would also exist in UK waters although, it is likely to be regionally significant (North Sea and Celtic and Irish Seas) during operations. Close proximity to noise created by geotechnical activity also has potential to cause injury, although evidence for the impact and level of risk is limited. This is also mitigated through guidance on operations such as soft start and on board marine mammal observers.</p>
<p>9.5: List of main conservation measures</p>	<p>MC02 Adapt/manage exploitation of energy resources. Guidance for the protection of marine European Protected Species from deliberate injury, killing and disturbance has been drafted (JNCC 2010a; Marine Scotland, 2014). Marine Industries generate a variety of noise through activities such as geophysical surveys (e.g. seismic surveys (JNCC 2017)), construction (e.g. pile driving (JNCC 2010b)) and decommissioning (e.g. use of explosives (2010c)). As part of the licencing procedures, developers and operators are required to utilise JNCC guidelines to minimise the risk of injury to cetaceans when undertaking such activities (JNCC, 2010b, 2010c; 2017; 2023; 2025; JNCC, Natural England & Cefas, 2025). The guidelines advise on conducting marine mammal observations prior to and during the activity and, where suitable, utilising procedures such as soft start (gradual</p>

introduction of the sound) to reduce and avoid direct harm to animals. A review of the marine mammal observer data demonstrated the effectiveness of soft start approach (Stone et al., 2017).

9.5: List of main conservation measures

MG05 Reduce bycatch and incidental killing of non-target species: The UK is implementing the EU Technical Conservation Measures Regulation transposed into UK regulations which lays down measures concerning incidental catches of vulnerable species in fisheries, and more generally the bycatch obligations within the Habitats Regulations. Since 2004, a dedicated bycatch monitoring programme has been in place, with both dedicated and non-dedicated onboard observers collecting data on bycatch numbers. These data inform implementation and potential effectiveness of measures such as pingers. There is a requirement for all fishing vessels over 12m using gill nets or entanglement nets to use pingers under the criteria laid out in the regulation. Inshore Vessel Monitoring System (iVMS) devices are being implemented for under-12 metre fishing vessels, allowing data on latitude, longitude, course and speed to be recorded and help improve the management and sustainability of the marine environment. Legislation to make iVMS mandatory on under-12 metre vessels is expected to come into effect in 2024 in England. In Scotland, consultation on the introduction mandatory electronic tracking for under-12 metre vessels was carried out in late 2023. Legislation requiring iVMS for under-12 metre vessels operating in Welsh waters has been in place since 2022. Since February 2022 it has been mandatory for under-10 metre fishing vessels in English and Welsh waters to create and submit a catch record for every fishing trip through the Catch Recording Application (Catch App or Record your Catch). Data is collected on vessel, trip, gear, area fished and catch and can be used to inform on fishing activity by gear type and species. Furthermore, the UK Marine Wildlife Bycatch Mitigation Initiative (published August 2022) aims to improve our understanding of bycatch and entanglement of sensitive marine species

	<p>through monitoring and scientific research, identify 'hotspot' or high-risk areas/gear types/fisheries in which to focus monitoring and mitigation, and develop and implement effective measures to minimise bycatch/entanglement. Currently work is progressing towards development of a bycatch risk framework across all PET species to apply all available evidence and support targeted monitoring.</p>
9.5: List of main conservation measures	<p>MG04 Control/eradication of illegal killing, fishing and harvesting. The Habitats Directive is transposed into UK law under the Habitat Regulations (HR) for England and Wales (as amended) and the Offshore Marine Conservation (Natural Habitats, &c.) Regulations 2007 (as amended), which make it an offence to kill, injure, capture or disturb European marine protected species. Similar legislation exists for Scottish and Northern Irish inshore waters.</p>
9.5: List of main conservation measures	<p>MH01 Reduce impact of military installations and activities. To reduce the risk of noise impact on marine mammals, the UK Ministry of Defence (MOD) has a Statement of Intent with UK Statutory Nature Conservation Bodies concerning conduct in relation to marine disturbance. The MOD has developed a real-time alert procedure for naval training operations. This enables localised information on cetacean sightings to be incorporated into the training schedule and for operations to be relocated if necessary.</p>
9.5: List of main conservation measures	<p>MK01 Reduce impact of mixed source pollution. The impact of chemical pollution on Risso's dolphin remains an issue (Jepson et al, 2016), however, establishing measures beyond the historic ban on PCB use, has not been achieved to date. Further information is required to understand where exposure is occurring to be able to identify appropriate measures.</p>
9.5: List of main conservation measures	<p>MC03 Adapt/manage renewable energy installation, facilities and operation (excl. hydropower and abstraction activities). Guidance for the protection of marine European Protected Species from deliberate injury, killing and</p>

disturbance has been drafted (JNCC 2010a; Marine Scotland, 2014). Marine Industries generate a variety of noise through activities such as geophysical surveys (e.g. seismic surveys (JNCC 2017)), construction (e.g. pile driving (JNCC 2010b)) and decommissioning (e.g. use of explosives (2010c)). As part of the licencing procedures, developers and operators are required to utilise JNCC guidelines to minimise the risk of injury to cetaceans when undertaking such activities (JNCC, 2010b, 2010c; 2017; 2023; 2025; JNCC, Natural England & Cefas, 2025). The guidelines advise on conducting marine mammal observations prior to and during the activity and, where suitable, utilising procedures such as soft start (gradual introduction of the sound) to reduce and avoid direct harm to animals. A review of the marine mammal observer data demonstrated the effectiveness of soft start approach (Stone et al., 2017).

9.5: List of main conservation measures

MG01 Management of professional/commercial fishing, shellfish and seaweed harvesting (incl. restoration of habitats). Fisheries Management Plans (FMPs) are currently being developed across all administrations for fisheries with perceived threats or pressures to the marine environment. FMPs are required under the Fisheries Act 2020 which provides the framework for management fisheries outside the EU Common Fisheries Policy. The Joint Fisheries Statement (agreeing the delivery of the 8 objectives of the Fisheries Act 2020) sets out plans for 43 FMPs. Publication of FMPs started last year and is expected to continue for 2-3 years. Some are being jointly developed, others by a single authority for its own waters. 6 FMPs have now been published.