

Reporting under the Habitat Regulations (as amended)¹

2019-2024

Conservation status assessment for the species:

S2032 - White-beaked dolphin

(*Lagenorhynchus albirostris*)

United Kingdom



Scottish Government
Riaghaltas na h-Alba
gov.scot



Cyfoeth
Naturiol
Cymru
Natural
Resources
Wales



Department of
Agriculture, Environment
and Rural Affairs

An Roinn
Talmhaíochta, Comhshaoil
agus Gnóthaí Tuaithe

Department of
Fairmin, Environment
an' Kintra Matters

www.daera-ni.gov.uk



Northern Ireland Environment Agency
Gníomhaireacht Comhshaoil Thuaisceart Éireann
www.daera-ni.gov.uk | www.niea.gov.uk | www.norlin-airian-environment.gov.uk

¹ Habitat Regulations (as amended):

- The Conservation of Habitats and Species Regulations 2017 (as amended), Regulation 9A
- The Conservation of Offshore Marine Habitats and Species Regulations 2017 (as amended), Regulation 6A
- Report under The Conservation (Natural Habitats, &c.) Regulations 1994 (as amended), regulation 3ZA
- The Conservation (Natural Habitats, etc.) Regulations (Northern Ireland) 1995 (as amended), regulation 3ZA

For further information please contact:

Joint Nature Conservation Committee. Quay House, 2 East Station Road, Fletton Quays, Peterborough, PE2 8YY. <https://jncc.gov.uk>

This report was produced by JNCC in collaboration with the UK Country Nature Conservation Bodies (CNCBs) and country governments.

This document should be cited as:

JNCC, Department of Agriculture, Environment and Rural Affairs, Natural England, Natural Resources Wales & NatureScot. (2026). Conservation status assessment for the species: S2032 White-beaked dolphin (*Lagenorhynchus albirostris*).

This resource and any accompanying material (e.g. maps, data, images) is published by JNCC under the [Open Government Licence](#) (OGLv3.0 for public sector information), unless otherwise stated. Note that some images (maps, tables) may not be copyright JNCC; please check sources for conditions of re-use.

The views and recommendations presented in this resource do not necessarily reflect the views and policies of JNCC.

Important note - Please read

- The information in this document represents the United Kingdom Reporting under the Habitat Regulations (as amended)¹, for the period 2019-2024.
- It is based on supporting information provided by Joint Nature Conservation Committee and UK Country Nature Conservation Bodies (CNCBs), which is documented separately.
- The Habitats Regulations reporting 2019-2024 Approach Document provides details on how this supporting information contributed to the UK Report and the fields that were completed for each parameter.
- Map showing the distribution and range of the species is included.
- Explanatory notes (where provided) are included at the end. These provide additional audit trail information to that included within the assessments. Further underpinning explanatory notes are available in the related country reports.
- Some of the reporting fields have been left blank because either: (i) there was insufficient information to complete the field; (ii) completion of the field was not obligatory; and/or (iii) the field was not relevant to this species (section 12 National Site Network coverage for Annex II species).

Further details on the approach to the Habitats Regulations Reporting 2019-2024 are available on the [JNCC website](#).

Assessment Summary: White-beaked dolphin

Distribution and Range Map

Distribution and Range
White-beaked Dolphin




Figure 1: United Kingdom distribution and range map for S2032 - White-beaked dolphin (*Lagenorhynchus albirostris*). The 50km grid square distribution map is based on available species records within the current reporting period.

Table 1: Table summarising the conservation status for S2032 - White-beaked dolphin (*Lagenorhynchus albirostris*). Overall conservation status for species is based on assessments of range, population, habitat for the species, and future prospects.

Overall Conservation Status (see section 11)

Unknown (XX)

Breakdown of Overall Conservation Status

Range (see section 5)

Favourable (FV)

Population (see section 6)

Favourable (FV)

Habitat for the species (see section 7)

Unknown (XX)

Future prospects (see section 10)

Unknown (XX)

List of Sections

National Level	5
1. General information	5
2. Maps	5
3. Information related to Annex V Species	5
Biogeographical Level	7
4. Biogeographical and marine regions	7
5. Range	7
6. Population	9
7. Habitat for the species	12
8. Main pressures	13
9. Conservation measures	14
10. Future prospects	16
11. Conclusions	16
12. UK National Site Network (pSCIs, SCIs, SACs) coverage for Annex II species	18
13. Complementary information	19
14. References	20
Biogeographical and marine regions	20
Main pressures	23
15. Explanatory Notes	25

National Level

1. General information

1.1 Country	United Kingdom
1.2 Species code	S2032
1.3 Species scientific name	<i>Lagenorhynchus albirostris</i>
1.4 Alternative species scientific name	
1.5 Common name	White-beaked dolphin
Annex(es)	IV

2. Maps

2.1 Sensitive species	No
2.2 Year or period	2019-2022
2.3 Distribution map	Yes
2.4 Distribution map; Method used	Complete survey or a statistically robust estimate

2.5 Additional information

The distribution map is based on verified sightings data of white-beaked dolphin between 2019 and 2024. The sightings were collated from SCANS IV, Pelagis French surveys, NBN Atlas, European Seabirds at Sea, the Joint Cetacean Data Programme, POSEIDON project, University of Aberdeen, The Crown Estate Marine Data Exchange, Whale and Dolphin Conservation, Hebridean Whale and Dolphin Trust, ORCA, Sea Watch Foundation, Marine Discovery Penzance, Sussex Dolphin Project, Cornwall Seal Group Research Trust, Cardigan Bay Marine Wildlife Centre, Natural England, Sea Trust and The Royal Society for the Protection of Birds (RSPB).

3. Information related to Annex V Species

3.1 Is the species taken in the wild / exploited?

3.2 What measures have been taken?

a) Regulations regarding access to property

b) Temporary or local prohibition on the taking of specimens in the wild and exploitation

c) Regulation of the periods and/or methods of taking specimens

d) Application of hunting and fishing rules which take account of the conservation of such populations

e) Establishment of a system of licences for taking specimens or of quotas

f) Regulation of the purchase, sale, offering for sale, keeping for sale, or transport for sale of specimens

g) Breeding in captivity of animal species as well as artificial propagation of plant species

Other measures

Other measures description

3.3: Hunting bag or quantity taken in the wild for Mammals and Acipenseridae (Fish)

a) Unit

Table 2: Quantity taken from the wild during the reporting period (see 3.3a for units). For species with defined hunting seasons, Season 1 refers to 2018/2019 (autumn 2018 to spring 2019), and Season 6 to 2023/2024. For species without hunting seasons, data are reported by calendar year: Year 1 is 2019, and Year 6 is 2024.

	Season/ year 1	Season/ year 2	Season/ year 3	Season/ year 4	Season/ year 5	Season/ year 6
b) Minimum	-	-	-	-	-	-
c) Maximum	-	-	-	-	-	-
d) Unknown	-	-	-	-	-	-

3.4: Hunting bag or quantity taken in the wild; Method used

3.5: Additional information

No additional information

Biogeographical Level

4. Biogeographical and marine regions

4.1 Biogeographical or marine region where the species occurs MATL

4.2 Sources of information

See section 14 References

5. Range

5.1 Surface area (km²) 488,645

5.2 Short-term trend; Period 2013-2024

5.3 Short-term trend; Direction Stable

5.4 Short-term trend; Magnitude

a) Estimated minimum

b) Estimated maximum

c) Pre-defined range

d) Unknown

e) Type of estimate

f) Rate of decrease

5.5 Short-term trend; Method used Complete survey or a statistically robust estimate used

5.6 Long-term trend; Period 1994-2024

5.7 Long-term trend; Direction Stable

**5.8 Long-term trend;
Magnitude****a) Minimum****b) Maximum****c) Rate of decrease**

5.9 Long-term trend; Method used Complete survey or a statistically robust estimate used**5.10 Favourable Reference Range (FRR)****a) Area (km²)** 475,589**b) Pre-defined increment****c) Unknown** No**d) Method used** Model-based approach**e) Quality of information** moderate**5.11 Change and reason for change in surface area of range****a) Change** Yes**b) Genuine change** No**c) Improved knowledge or more accurate data** Yes**d) Different method** No**e) No information** No**f) Other reason** No**g) Main reason** Improved knowledge/more accurate data**5.12 Additional information**

The distribution is based on verified sightings data of white-beaked dolphin between 2019 and 2024. The sightings were collated from SCANS IV, Pelagis French surveys, NBN Atlas, European Seabirds at Sea, the Joint Cetacean Data Programme, POSEIDON project, University of Aberdeen, The Crown Estate Marine Data Exchange, Whale and Dolphin Conservation, Hebridean Whale and Dolphin Trust, ORCA, Sea Watch Foundation, Marine Discovery Penzance, Sussex Dolphin Project, Cornwall Seal

Group Research Trust, Cardigan Bay Marine Wildlife Centre, Natural England, Sea Trust and The Royal Society for the Protection of Birds (RSPB).

UK waters represent the southern range limit for this species (Williamson et al. 2023), therefore white-beaked dolphin is restricted to northern waters of the UK EEZ, the North Sea and an isolated population found in the western Channel. Over recent years, white-beaked dolphin have been reported in the eastern Channel and the northern Irish sea though these are less common. However, the range has been extended to include these areas.

The FRR was based on an analysis of effort-related survey spanning 1994-2010 compiled for the Joint Cetacean Protocol (JCP) undertaken by Paxton et al. (2016). The estimated range was based on a modelled prediction of white-beaked dolphin distribution during August 2010 and adapted based on additional sightings data and expert knowledge (see Paxton et al., 2016 for further detail).

Since the 2019 Habitats Directive Article 17 assessments, the FRR has changed due to the removal of the EEZ extension into offshore waters west of Scotland. This area has been removed due to lack of data for all species, and subsequent impact on confidence in assessments. This does not represent genuine change in FRR.

6. Population

6.1 Year or period 2022

6.2 Population size (in reporting unit)

a) Unit number of individuals

b) Minimum 29,484

c) Maximum 75,536

d) Best single value 47,192

6.3 Type of estimate 95% confidence interval

6.4 Quality of extrapolation to reporting unit high

6.5 Additional population size (using population unit other than reporting unit)

a) Unit

b) Minimum

c) Maximum

d) Best single value

e) Type of estimate

6.6 Population size; Method used Complete survey or a statistically robust estimate used

6.7 Short-term trend; Period 2016-2022

6.8 Short-term trend; Direction Increasing

6.9 Short-term trend; Magnitude

a) Estimated minimum

b) Estimated maximum

c) Pre-defined range Increasing 13 - 25%

d) Unknown No

e) Type of estimate 95% confidence interval

f) Rate of decrease

6.10 Short-term trend; Method used Complete survey or a statistically robust estimate used

6.11 Long-term trend; Period 2005-2022

6.12 Long-term trend; Direction Increasing

6.13 Long-term trend; Magnitude

a) Minimum

b) Maximum

c) Confidence interval

d) Rate of decrease

6.14 Long-term trend; Method used Complete survey or a statistically robust estimate used

6.15 Favourable Reference Population (FRP)

ai) Population size	21,256
aii) Unit	number of individuals
b) Pre-defined increment	
c) Unknown	No
d) Method used	Model-based approach
e) Quality of information	high

6.16 Change and reason for change in population size

a) Change	Yes
b) Genuine change	Yes
c) Improved knowledge or more accurate data	No
d) Different method	No
e) No information	No
f) Other reason	No
g) Main reason	Genuine change

6.17 Additional information

The abundance of white-beaked dolphins within the UK EEZ has continued to increase since 2005 (Hammond et al., 2021; Gilles, et al., 2023). This follows the same pattern of increase for the wider SCANS survey. The abundance of white-beaked dolphin has increased in the NE Atlantic based on the latest SCANS survey in 2022 in comparison to previous surveys in 2015 and 2005 (Gilles et al. 2023). In previous abundance estimates the confidence intervals overlap considerably indicating that there is no significant difference between these estimates.

The wider context for white-beaked dolphin population in the NE Atlantic depends on the outputs from NASS 2024 survey in the NAMMCO region once published.

The FRV for population (21256 CV: 0.293 CI: 12117-37287) was calculated based on estimates from SCANS II in 2005 (Hammond, et al., 2021) and CODA in 2007 (Hammond, et al., 2009), supplemented with density estimates from neighbouring regions to fill data gaps within the UK EEZ and limit extrapolation where possible; ObSERVE in Irish waters (Rogan, e al., 2018), and NASS and T-NASS (Pike, et al.,

2019a; Pike, et al., 2019b) and NILS (Leonard and Øien, 2020a; Leonard and Øien, 2020b) surveys in the NAMMCO region.

Since the 2019 Habitats Directive Article 17 assessments, the FRV has changed due to the removal of the EEZ extension into offshore waters west of Scotland. This area has been removed due to lack of data for all species, and subsequent impact on confidence in assessments. This does not represent genuine change in FRV.

As short-term trend in population has been determined as increasing, the pre-defined range field has been used to indicate a magnitude. However, population estimates have been calculated from large-scale surveys at approximately decadal intervals, and the confidence intervals associated with population estimates are wide. Thus, confidence in the magnitude is low.

6.18 Age structure, mortality and reproduction deviation Unknown

7. Habitat for the species

7.1 Sufficiency of area and quality of occupied habitat (for long-term survival)

a) Is area of occupied habitat sufficient? Unknown

b) Is quality of occupied habitat sufficient? Unknown

c) If No or Unknown, is there a sufficiently large area of unoccupied habitat of suitable quality? Unknown

7.2 Sufficiency of area and quality of occupied habitat; Method used

a) Sufficiency of area of occupied habitat; Method used Based mainly on expert opinion with very limited data

b) Sufficiency of quality of occupied habitat; Method used Based mainly on expert opinion with very limited data

7.3 Short-term trend; Period

7.4 Short-term trend; Direction Unknown

7.5 Short-term trend; Method used	Based mainly on expert opinion with very limited data
7.6 Long-term trend; Period	
7.7 Long-term trend; Direction	Unknown
7.8 Long-term trend; Method used	Based mainly on expert opinion with very limited data

7.9 Additional information

Direct evidence of cetacean habitat quality is limited as presently, a comprehensive understanding of the key elements important to the species is undetermined. In some cases, conclusions for species range and population could be indicative of habitat quality by proxy, however confidence in assessment outputs would be low.

8. Main pressures

8.1 Characterisation of pressures

Table 3: Pressures affecting the species, including timing and importance/impact ranking. Pressures are defined as factors acting currently and/or during the reporting period (2019–2024). Rankings are: High (direct/immediate influence and/or large spatial extent) and Medium (moderate direct/immediate influence, mainly indirect and/or regional extent).

Pressure	Timing	Ranking
Geotechnical surveying	Ongoing and likely to be in the future	Medium (M)
Residential, commercial and industrial activities and structures generating noise, light, heat or other forms of pollution	Ongoing and likely to be in the future	Medium (M)
Marine fish and shellfish harvesting causing reduction of species/prey populations and disturbance of species (professional)	Ongoing and likely to be in the future	Medium (M)
Bycatch and incidental killing (due to fishing and hunting activities)	Ongoing and likely to be in the future	Medium (M)
Mixed source marine water pollution (marine and coastal)	Ongoing and likely to be in the future	Medium (M)
Change of species distribution (natural newcomers) due to climate change	Ongoing and likely to be in the future	Medium (M)

Decline or extinction of related species (e.g. food source / prey, predator / parasite, symbiote, etc.) due to climate change	Ongoing and likely to be in the future	Medium (M)
Threats and pressures from outside the Member State	Ongoing and likely to be in the future	Medium (M)

8.2 Sources of information

See section 14 References

8.3 Additional information

PC07: Regional pressure in the North Sea and the Irish Sea.

PJ13: Pressure PJ13 is the closest appropriate pressure but is being used to consider all changes in species distribution, including range contractions (i.e., not only natural newcomers).

PX02: Relating to continued whaling of this species outside of UK waters which may be having an impact on populations.

9. Conservation measures

9.1: Status of measures

a) Are measures needed?

Yes

b) Indicate the status of measures

Measures identified and taken

9.2 Main purpose of the measures taken

Maintain the current range, population and/or habitat for the species

9.3 Location of the measures taken

Both inside and outside National Site Network

9.4 Response to measures

Medium-term results (within the next two reporting periods, 2025–2036)

9.5 List of main conservation measures

Table 4: Key conservation measures addressing current pressures and/or anticipated threats during the next two reporting periods (2025–2036). Measures are ranked by importance/impact: High (direct/immediate influence and/or large spatial extent) and Medium (moderate direct/immediate influence, mainly indirect and/or regional extent).

Conservation measure	Ranking
Adapt/manage exploitation of energy resources	High (H)
Adapt/manage renewable energy installation, facilities and operation (excl. hydropower and abstraction activities)	High (H)
Management of professional/commercial fishing, shellfish and seaweed harvesting (incl. restoration of habitats)	High (H)
Reduce bycatch and incidental killing of non-target species	High (H)
Reduce impact of mixed source pollution	High (H)
Control/eradication of illegal killing, fishing and harvesting of wild plants, fungi and animals	High (H)
Reduce impact of military installations and activities	High (H)

9.6 Additional information

This species is not an Annex II species and therefore the designation of SACs is not required, as stipulated in the Habitats Regulations. However, as a European Protected Species, protection is provided throughout UK waters and it is an offence to kill, injure or disturb. The UK remains committed to the conservation of marine mammals in UK waters and the implementation of measures to mitigate the impact of pressures and conservation measures have been undertaken in the UK and adjacent waters as part of the requirements of the Habitats Regulations. Such measures include monitoring bycatch, monitoring strandings data to monitor current and identify emerging pressures, application of appropriate management measures, and noise monitoring and mitigation with regards to offshore industry. This is reflected in the list of conservation measures under field 9.5. The UK also supports a range of international agreements and conventions on the conservation of marine mammals and the marine environment. For example: The Convention on Migratory Species; the Convention for the Protection of the Marine Environment of the North-East Atlantic (OSPAR). A UK Cetacean Conservation Strategy is currently in development, due for publication shortly. The strategy is intended to support decision making and identify actions necessary to maintain or improve the conservation status of cetaceans in UK waters. Defra and devolved administrations fund national strandings schemes for cetaceans which aim to: collate, analyse and report data for all cetacean strandings around the coast of the UK; determine the causes of death (both natural and anthropogenic) in stranded cetaceans, including bycatch and physical trauma and; undertake surveillance on the incidence of disease in stranded cetaceans in order to identify any substantial new threats to their conservation status.

10. Future prospects

10.1a Future trends of parameters

ai) Range	Unknown
-----------	---------

bi) Population	Unknown
----------------	---------

ci) Habitat for the species	Unknown
-----------------------------	---------

10.1b Future prospects of parameters

aii) Range	Unknown
------------	---------

bii) Population	Unknown
-----------------	---------

cii) Habitat for the species	Unknown
------------------------------	---------

10.2 Additional information

Uncertainty of the prospective impact of climate change on the range and distribution of this species in UK waters is driving the 'Unknown' outcomes for Future Prospects (Williamson et al., 2021).

11. Conclusions

11.1 Range	Favourable (FV)
------------	-----------------

11.2 Population	Favourable (FV)
-----------------	-----------------

11.3 Habitat for the species	Unknown (XX)
------------------------------	--------------

11.4 Future prospects	Unknown (XX)
-----------------------	--------------

11.5 Overall assessment of Conservation Status	Unknown (XX)
--	--------------

11.6 Overall trend in Conservation Status	Stable
---	--------

11.7 Change and reason for change in conservation status

a) Change	No
-----------	----

b) Genuine change	
-------------------	--

c) Improved knowledge or more accurate data

d) Different method

e) No information

f) Other reason

g) Main reason

11.7 Change and reason for change in conservation status trend

a) Change Yes

b) Genuine change No

c) Improved knowledge or more accurate data Yes

d) Different method No

e) No information No

f) Other reason No

g) Main reason Improved knowledge/more accurate data

11.8 Additional information

Conclusion on Range reached because: (i) the short-term trend direction in Range surface area is stable and (ii) the current Range surface area is larger than the Favourable Reference Range.

Conclusion on Population reached because: (i) the short-term trend direction in Population size is increasing; and (ii) the best estimate for population size is greater than the Favourable Reference Population.

Conclusion on Habitat for the species reached because: (i) it is unknown whether the area of habitat is sufficiently large; (ii) it is unknown if habitat quality is sufficient for the long-term survival of the species; and (iii) the short-term trend in area and quality of habitat is unknown.

Conclusion on Future prospects reached because: (i) the Future prospects for Range are Unknown; (ii) the Future prospects for Population are Unknown; and (iii) the Future prospects for Habitat for the species are Unknown.

Overall assessment of Conservation Status is Unknown because all conclusions are Unknown.

Overall trend in Conservation Status is based on the combination of the short-term trends for Range - stable, Population - increasing, and Habitat for the species - unknown.

Evidence suggests trends in distribution are becoming apparent for this species likely driven by climate change, which is contributing to some uncertainty in conclusions for future prospects for this species.

12. UK National Site Network (pSCIs, SCIs, SACs) coverage for Annex II species

12.1 Population size inside the pSCIs, SCIs and SACs network

a) Unit

b) Minimum

c) Maximum

d) Best single value

12.2 Type of estimate

12.3 Population size inside the network; Method used

12.4 Short-term trend of population size within the network; Direction

12.5 Short-term trend of population size within the network; Method used

12.6 Short-term trend of habitat for the species inside the pSCIs, SCIs and SACs network; Direction

12.7 Short-term trend of habitat for the species inside the pSCIs, SCIs and SACs network; Method used

12.8 Additional information

No additional information

13. Complementary information

13.1 Justification of percentage thresholds for trends

No justification information

13.2 Trans-boundary assessment

No trans-boundary assessment information

13.2 Other relevant information

No other relevant information

14. References

Biogeographical and marine regions

4.2 Sources of information

- Williamson, M.J., ten Doeschate, M.T.I., Deaville, R., Brownlow, A.C., Taylor, N.L., 2021. Cetaceans as sentinels for informing climate change policy in UK waters. *Marine Policy* 131, 104634. <https://doi.org/10.1016/j.marpol.2021.104634>
- Gilles, A, Authier, M, Ramirez-Martinez, NC, Araújo, H, Blanchard, A, Carlström, J, Eira, C, Dorémus, G, Fernández-Maldonado, C, Geelhoed, SCV, Kyhn, L, Laran, S, Nachtsheim, D, Panigada, S, Pigeault, R, Sequeira, M, Sveegaard, S, Taylor, NL, Owen, K, Saavedra, C, Vázquez-Bonales, JA, Unger, B, Hammond, PS (2023). Estimates of cetacean abundance in European Atlantic waters in summer 2022 from the SCANS-IV aerial and shipboard surveys. Final report published 29 September 2023. 64 pp. <https://www.tiho-hannover.de/itaw/scans-iv-survey>
- Paxton, C.G.M., Scott-Hayward, L., Mackenzie, M., Rexstad, E. & Thomas, L. (2016) Revised Phase III Data Analysis of Joint Cetacean Protocol Data Resource, JNCC Report No. 517, JNCC, Peterborough, ISSN 0963-8091. <https://hub.jncc.gov.uk/assets/01adfabd-e75f-48ba-9643-2d594983201e>
- Pike, D.G., Gunnlaugsson, T., Mikkelsen, B., Halldórsson, S.D. & Víkingsson, G.A. (2019a). Estimates of the Abundance of Cetaceans in the Central North Atlantic Based on the NASS Icelandic and Faroese Shipboard Surveys Conducted in 2015. NAMMCO Scientific Publications 11. <https://doi.org/10.7557/3.4941>
- Leonard, D. M. & Øien, N. I. (2020a). Estimated Abundances of Cetacean Species in the Northeast Atlantic from Norwegian Shipboard Surveys Conducted in 2014–2018. NAMMCO Scientific Publications 11. <https://doi.org/10.7557/3.4694>
- Brownlow, A., Davison, N. and ten Doeschate, M. 2019. Scottish Marine Animal Stranding Scheme (SMASS) Annual Report 2018. Available at: https://strandings.org/wp-content/uploads/2021/05/SMASS_Annual_Report_2019.pdf [Accessed 07 Nov 2024]
- Deaville, R. (compiler). 2019:2024. Annual Reports for the period 1st January to 31st December. UK Cetacean Strandings Investigation Programme (CSIP).
- Fall, J. and Skern-Mauritzen, M., 2014. White-beaked dolphin distribution and association with prey in the Barents Sea. *Marine Biology Research*, 10(10), pp.957-971.

Jansen, O.E., 2013. Fishing for food: feeding ecology of harbour porpoises *Phocoena phocoena* and white-beaked dolphins *Lagenorhynchus albirostris* in Dutch waters. Wageningen University and Research.

Jansen, O.E., Leopold, M.F., Meesters, E.H. and Smeenk, C., 2010. Are white-beaked dolphins *Lagenorhynchus albirostris* food specialists? Their diet in the southern North Sea. *Journal of the Marine Biological Association of the United Kingdom*, 90(8), pp.1501-1508.

Jepson, P.D., Deaville, R., Acevedo-Whitehouse, K., Barnett, J., Brownlow, A., Brownell Jr, R.L., Clare, F.C., Davison, N., Law, R.J., Loveridge, J. and Macgregor, S.K., 2013. What caused the UK's largest common dolphin (*Delphinus delphis*) mass stranding event?. *PLoS One*, 8(4), p.e60953.

JNCC. 2010a. The protection of marine European Protected Species from deliberate injury, killing and disturbance. Guidance for the marine area in England and Wales and the UK offshore marine area. Available on request from JNCC.

JNCC. 2010b. Statutory nature conservation agency protocol for minimising the risk of injury to marine mammals from Piling noise. 2010. JNCC Peterborough. United Kingdom. Available at: <https://data.jncc.gov.uk/data/31662b6a-19ed-4918-9fab-8fbcff752046/JNCC-CNCB-Piling-protocol-August2010-Web.pdf> [Accessed 06 Nov 2024]

JNCC. 2010c. JNCC guidelines for minimising the risk of injury to marine mammals from using explosives. August 2010. Available at: <https://data.jncc.gov.uk/data/24cc180d-4030-49dd-8977-a04ebe0d7aca/JNCC-Guidelines-Explosives-Guidelines-201008-Web.pdf> [Accessed 06 Nov 2024]

JNCC. 2017. JNCC guidelines for minimising the risk of injury to marine mammals from geophysical surveys. Available at: <https://data.jncc.gov.uk/data/e2a46de5-43d4-43f0-b296-c62134397ce4/jncc-guidelines-seismicsurvey-aug2017-web.pdf> [Accessed 06 Nov 2024]

JNCC. 2023. JNCC guidance for the use of Passive Acoustic Monitoring in UK waters for minimising the risk of injury to marine mammals from offshore activities. JNCC, Peterborough. Available at: <https://hub.jncc.gov.uk/assets/fb7d345b-ec24-4c60-aba2-894e50375e33> [Accessed 06 Nov 2024]

Malinauskaite, L., Cook, D., Davíðsdóttir, B., Karami, M.P., Koenigk, T., Kruschke, T., Ögmundardóttir, H. and Rasmussen, M., 2022. Connecting the dots: An interdisciplinary perspective on climate change effects on whales and whale watching in Skjalfandi Bay, Iceland. *Ocean & Coastal Management*, 226, p.106274.

Mann, D., Hill-Cook, M., Manire, C., Greenhow, D., Montie, E., Powell, J., Wells, R., Bauer, G., Cunningham-Smith, P., Lingenfelter, R. and DiGiovanni Jr, R., 2010. Hearing loss in stranded odontocete dolphins and whales. *PLoS One*, 5(11), p.e13824.

Marine Scotland. 2014. The protection of Marine European Protected Species from injury and disturbance. Guidance for Scottish Inshore Waters.

Megson, D., Brown, T., Jones, G.R., Robson, M., Johnson, G.W., Tiktak, G.P., Sandau, C.D. and Reiner, E.J., 2022. Polychlorinated biphenyl (PCB) concentrations and profiles in marine mammals from the North Atlantic Ocean. *Chemosphere*, 288, p.132639.

Merchant, N.D., Pirotta, E., Barton, T.R. and Thompson, P.M., 2014. Monitoring ship noise to assess the impact of coastal developments on marine mammals. *Marine Pollution Bulletin*, 78(1-2), pp.85-95.

Middel, H. and Verones, F., 2017. Making marine noise pollution impacts heard: the case of cetaceans in the North sea within life cycle impact assessment. *Sustainability*, 9(7), p.1138.

Morizur, Y., Berrow, S.D., Tregenza, N.J.C., Couperus, A.S. and Pouvreau, S., 1999. Incidental catches of marine-mammals in pelagic trawl fisheries of the northeast Atlantic. *Fisheries Research*, 41(3), pp.297-307.

Nachtigall, P.E., Mooney, T.A., Taylor, K.A., Miller, L.A., Rasmussen, M.H., Akamatsu, T., Teilmann, J., Linnenschmidt, M. and Vikingsson, G.A., 2008. Shipboard measurements of the hearing of the white-beaked dolphin *Lagenorhynchus albirostris*. *Journal of Experimental Biology*, 211(4), pp.642-647.

Pike, D.G., Gunnlaugsson, T., Sigurjónsson, J. and Vikingsson, G., 2019. Distribution and abundance of cetaceans in Icelandic waters over 30 years of aerial surveys. *NAMMCO Scientific Publications*, 11.

Plint, T., ten Doeschate, M.T., Brownlow, A.C., Davison, N.J., Hantke, G., Kitchener, A.C., Longstaffe, F.J., McGill, R.A., Simon-Nutbrown, C. and Magill, C.R., 2023. Stable isotope ecology and interspecific dietary overlap among dolphins in the Northeast Atlantic. *Frontiers in Marine Science*, 10, p.1111295.

Reverberi, M., 2023. The non-silent world: acoustic responses of white-beaked dolphins (*Lagenorhynchus albirostris*) to changes in maritime traffic: a case study during the covid-19 anthropause in Skjálfandi Bay, Iceland (Doctoral dissertation). Available at: https://skemman.is/bitstream/1946/45856/1/CMMThesis_MathieuReverberi.pdf [Accessed 07 Nov 2024]

Scottish Marine Animal Stranding Scheme. 2022. Scottish Marine Animal Stranding Scheme (SMASS) Annual Report 2021. Available: <https://strandings.org/wp-content/uploads/2024/06/SMASS-Annual-Report-2021-final.pdf> [Accessed 07 Nov 2024]

Simmonds, M., McLellan, F., Entrup, N., & Nunny, L. (2021). Whaling in Europe: An Ongoing Welfare and Conservation Concern In: Under Pressure: The need to protect whales and dolphins in European waters. An OceanCare Report. Available at: https://www.oceancare.org/wp-content/uploads/2022/11/Animal_Species_Protection_Under-Pressure_Whales-and-Dolphins_EU_Report_OceanCare_EN_146p_2021.pdf

[Accessed 06 Nov 2024]

Stone, C., Hall, K., Mendes, S. and Tasker, M. 2017. The effects of seismic operations in UK waters: analysis of Marine Mammal Observer data. *J. Cetacean Res. Manage.*, 16, pp.71-85.

Stone, C.J. 2003. The effects of seismic activity on marine mammals in UK waters, 1998-2000. JNCC Report No. 323. Available at: <https://data.jncc.gov.uk/data/bf3ea885-e5c5-4088-956b-4f5ff9ca0b56/JNCC-Report-323-FINAL-WEB.pdf> [Accessed 06 Nov 2024]

Stone, C.J. 2015. Implementation of and considerations for revisions to the JNCC guidelines for seismic surveys. JNCC Report No. 463b. Available at: <https://data.jncc.gov.uk/data/f7990481-7a99-414c-be04-b972da10c1b7/JNCC-Report-463b-FINAL-WEB.pdf> [Accessed 06 Nov 2024]

Stone, C.J. and Tasker, M.L. 2006. The effects of seismic airguns on cetaceans in UK waters. *J. Cetacean Res. Manage.*, 8(3), pp.255-263.

Van de Vijver, K.I., Hoff, P.T., Das, K., Van Dongen, W., Esmans, E.L., Jauniaux, T., Bouquegneau, J.M., Blust, R. and De Coen, W., 2003. Perfluorinated chemicals infiltrate ocean waters: link between exposure levels and stable isotope ratios in marine mammals. *Environmental science & technology*, 37(24), pp.5545-5550.

JNCC. 2025. JNCC guidelines for minimising the risk of injury to marine mammals from unexploded ordnance (UXO) clearance in the marine environment. JNCC, Aberdeen.

JNCC, Natural England and Cefas. 2025. JNCC, Natural England and Cefas position on the use of quieter piling methods and noise abatement systems when installing offshore wind turbine foundations. JNCC, Aberdeen.

Evans, P.G.H. and Waggitt, J.J. 2023. Modelled Distribution and Abundance of Cetaceans and Seabirds in Wales and Surrounding Waters. NRW Evidence Report, Report No: 646,

354 pp. Natural Resources Wales, Bangor.

Main pressures

8.2 Sources of information

No sources of information

15. Explanatory Notes

Field label	Note
8.1: Characterisation of pressures	<p>PJ12 Decline or extinction of related species (e.g. food source / prey, predator / parasite, symbiot, etc.) due to climate change. There is no current evidence for the effects of climate change on white-beaked dolphins. However, the effects of climate change on prey availability is likely to be mediated through variation in prey resource initially. Previous studies have highlighted that white-beaked dolphins may have a varied diet (Fall and Skern-Mauritzen, 2014; Jansen et al, 2010; Jansen, 2013) and are able to change distributions in response to changes in prey distribution (Malinauskaite et al., 2022; Pike et al., 2022). They may therefore adapt to new food sources, potentially reducing the impact of this threat. However, recent stable isotope analysis has highlighted that white-beaked dolphins showed low dietary plasticity, indicating limited ability to adapt to changes in prey availability (Plint et al., 2023). Further, a high overlap in diet preferences with short-beaked common dolphin was found (Plint et al., 2023). With suggestions that short-beaked common dolphin distributions are moving north (Williamson et al., 2021), and will overlap more with the distribution of white-beaked dolphin this may exacerbate the impact of this pressure.</p>
8.1: Characterisation of pressures	<p>PC07 Geotechnical surveying. The primary impact of this pressure is disturbance which is indirect with evidence of recovery/return once the pressure is removed. There is also a lesser risk of injury if in close proximity, which may indirectly lead to impairment and or mortality (Mann et al., 2010). White-beaked dolphins were the most frequently encountered species recorded during visual and/or acoustic observation between 1994 and 2010 in UK waters by Stone (2015) indicating relatively high exposure to seismic noise. A study in Iceland indicated white-beaked dolphins have noise sensitivity thresholds comparable to harbour porpoise and are therefore potentially vulnerable to high frequency activity (Nachtigall et al., 2008). Seismic and other</p>

geotechnical surveys have an immediate influence on white-beaked dolphins, causing disturbance. The species shows strong avoidance of seismic activity (Stone, 2015, Stone et al., 2017) with significant increases in fast swimming (Stone, 2003), orientation variation, and displaying strong lateral spatial avoidance (Stone and Tasker, 2006). Studies in Iceland have also demonstrated that more white-beaked dolphin whistles were detected during periods of reduced marine traffic (overnight and during Covid-19 pandemic), suggesting avoidance responses to vessel noise (Reverberi, 2023). This may indirectly influence survival and/or fecundity. Close proximity to noise created by geotechnical activity also has potential to cause injury, although evidence for the impact and level of risk is limited. This is also mitigated through guidance on operations such as soft start and on board marine mammal observers. Pressure is likely to be regionally significant during operations and will be higher in the North Sea and Celtic and Irish Seas.

8.1: Characterisation of pressures

PF12 Industrial or commercial activities and structures generating noise, light, heat or other forms of pollution. Cetaceans rely on echolocation for navigation, foraging and communication, making them sensitive to noise in the marine environment (Middel and Verones, 2017). Although various individual sources of disturbance have been identified as potential pressures in the pre-defined EU list, such as noise from shipping vessels or renewable energy devices, these pressures independently have not been identified as Medium or High risk to white-beaked dolphins in UK waters. The cumulative effect of such disturbances however, has the potential to increase energy expenditure through avoidance and decreased foraging. Furthermore, excessive anthropogenic noise may have the potential for auditory masking and reduced communication between individuals (Merchant et al., 2014). Studies in Iceland have also demonstrated that more white-beaked dolphin whistles were detected during periods of reduced marine traffic (overnight and during Covid-19 pandemic), suggesting avoidance responses to vessel noise (Reverberi, 2023).

However, there is currently a scarcity of evidence on the impact of anthropogenic noise on white-beaked dolphins.

8.1: Characterisation of pressures

PG01 Marine fish and shellfish harvesting (professional, recreational) causing reduction of species/prey populations and disturbance of species. Starvation was reported in 13% of animals examined through the through the UK Cetacean Stranding Investigation Programme (10 of 79 adults) between 2001-2017 (UK Cetacean Stranding Investigation Programme (CSIP) annual reports <http://ukstrandings.org/csip-reports>), although only one of 25 examined adult animals (4%) was thought to have died due to starvation between 2018 and 2022 (Deaville, 2021). White-beaked dolphins have a varied diet (Fall and Skern-Mauritzen, 2014) and are known to feed on at least 25 different fish species, many of which are commercially targeted by fisheries (e.g. whiting and cod) (Jansen et al, 2010; Jansen, 2013). However, recent stable isotope analysis has highlighted that white-beaked dolphins showed low dietary plasticity, indicating limited ability to adapt to changes in prey availability (Plint et al., 2023). Further, a high overlap in diet preferences with short-beaked common dolphin was found (Plint et al., 2023) and with suggestions that short-beaked common dolphin distributions are moving north (Williamson et al., 2021), overlapping more with the distribution of white-beaked dolphin the issue of prey depletion may be exacerbated. It should be noted that prey depletion can occur due to both natural and anthropogenic causes. No direct link has been established between commercial fishing practices and the cases of cetacean starvation recorded from the UK CSIP.

8.1: Characterisation of pressures

PG13 Bycatch and incidental killing (due to fishing and hunting activities). Bycatch in pelagic trawl nets in the North Sea has historically been noted as a risk for white-beaked dolphins (Morizur et al., 1999). Between 2000-2017, 3% (2 of 79 animals examined) of mortality investigated by the UK CSIP was caused by bycatch and there are two records (2011 - 2017) of white-beaked dolphin bycatch through the UK bycatch observer monitoring programme, one in pelagic

	<p>gear and one in a demersal set net. No stranded animals were determined as bycatch related between 2018 and 2022 (Deaville 2019:2024). Given the direct impact of bycatch, but with limited evidence of significant impact, the pressure is ranked as medium.</p>
8.1: Characterisation of pressures	<p>PK02 Mixed source marine water pollution (marine and coastal). The general impact of contaminants on cetaceans is well documented, including impacts on the immune system and reproduction (Jepson et al, 2016). The concentration is highly dependent on the age, sex, reproductive state and nutritional condition of the animals in addition to the intake via the food web. There is limited information on the severity of chemical pollutants found in white-beaked dolphins, however one study found that perfluorinated organochemicals (PFOCs) levels in white-beaked dolphins were amongst the highest in sampled marine mammal species stranded along the North Sea coast (Van De Vijver et al., 2003). Another study found concentrations of PCBs which exceeded toxic thresholds in 2 of 3 white-beaked dolphins sampled; one of these samples contained 155 different PCBs, the highest number found (jointly with Risso's; Megson et al., 2022).</p>
8.1: Characterisation of pressures	<p>PJ13 Change of species distribution (natural newcomers) due to climate change. There is limited evidence of this impact from around the UK but studies in Iceland have demonstrated a northward shift in this species due to environmental change and changes in the distribution/availability of prey (Malinauskaite et al., 2022; Pike et al., 2020).</p>
8.1: Characterisation of pressures	<p>PI04 Plant and animal disease, pathogens and pests. Necropsies of stranded animals highlights consistent evidence of parasitic infestation and infection from pathogens, with cause of death attributed to bacterial or parasitic infections in 24% of necropsied animals between (Deaville 2019:2024; Brownlow et al., 2019; Scottish Marine Animal Stranding Scheme, 2022).</p>
8.1: Characterisation of pressures	<p>PX02 Threats and pressures from outside the Member State. Related to the continued take of this species in</p>

neighbouring waters. White-beaked dolphins are hunted without quota in the Faroe Islands and Greenland although catch data is limited and are combined with Atlantic white-sided dolphin catch data (Simmonds et al., 2021), thus it is difficult to determine the extent of this pressure on white-beaked dolphin specifically. Numbers may be significant with over 400 white-sided/white-beaked dolphins recorded in some years over the past decade (<http://www.whaling.fo/en/regulated/450-years-of-statistics/catches>).

9.5: List of main conservation measures

MG05 Reduce bycatch and incidental killing of non-target species. The UK is implementing the EU Technical Conservation Measures Regulation transposed into UK regulations which lays down measures concerning incidental catches of vulnerable species in fisheries, and more generally the bycatch obligations within the Habitats Regulations. Since 2004, a dedicated bycatch monitoring programme has been in place, with both dedicated and non-dedicated onboard observers collecting data on bycatch numbers. These data inform implementation and potential effectiveness of measures such as pingers. There is a requirement for all fishing vessels over 12m using gill nets or entanglement nets to use pingers under the criteria laid out in the regulation. Inshore Vessel Monitoring System (iVMS) devices are being implemented for under-12 metre fishing vessels, allowing data on latitude, longitude, course and speed to be recorded and help improve the management and sustainability of the marine environment. Legislation to make iVMS mandatory on under-12 metre vessels is expected to come into effect in 2024 in England. In Scotland, consultation on the introduction mandatory electronic tracking for under-12 metre vessels was carried out in late 2023. Legislation requiring iVMS for under-12 metre vessels operating in Welsh waters has been in place since 2022. Since February 2022 it has been mandatory for under-10 metre fishing vessels in English and Welsh waters to create and submit a catch record for every fishing trip through the Catch Recording Application (Catch App or Record your Catch). Data is collected on vessel, trip, gear, area fished and catch and can be used to inform on fishing

	<p>activity by gear type and species. Furthermore, the UK Marine Wildlife Bycatch Mitigation Initiative (published August 2022) aims to improve our understanding of bycatch and entanglement of sensitive marine species through monitoring and scientific research, identify 'hotspot' or high-risk areas/gear types/fisheries in which to focus monitoring and mitigation, and develop and implement effective measures to minimise bycatch/entanglement. Currently work is progressing towards development of a bycatch risk framework across all PET species to apply all available evidence and support targeted monitoring.</p>
9.5: List of main conservation measures	<p>MK01 Reduce impact of mixed source pollution. The impact of chemical pollution on white-beaked dolphin remains an issue (Jepson et al, 2016), however, establishing measures beyond the historic ban on PCB use, has not been achieved to date. Further information is required to understand where exposure is occurring to be able to identify appropriate measures.</p>
9.5: List of main conservation measures	<p>MH01 Reduce impact of military installations and activities. To reduce the risk of noise impact on marine mammals, the UK Ministry of Defence (MOD) has a Statement of Intent with UK Statutory Nature Conservation Bodies concerning conduct in relation to marine disturbance. The MOD has developed a real-time alert procedure for naval training operations. This enables localised information on cetacean sightings to be incorporated into the training schedule and for operations to be relocated if necessary.</p>
9.5: List of main conservation measures	<p>MG04 Control/eradication of illegal killing, fishing and harvesting: The Habitats Directive is transposed into UK law under the Habitat Regulations (HR) for England and Wales (as amended) and the Offshore Marine Conservation (Natural Habitats, &c.) Regulations 2007 (as amended), which make it an offence to kill, injure, capture or disturb European marine protected species. Similar legislation exists for Scottish and Northern Irish inshore waters.</p>
9.5: List of main conservation measures	<p>MC02 Adapt/manage exploitation of energy resources: Guidance for the protection of marine European Protected Species from deliberate injury, killing and disturbance has</p>

been drafted (JNCC 2010a; Marine Scotland, 2014). Marine Industries generate a variety of noise through activities such as geophysical surveys (e.g. seismic surveys (JNCC 2017)), construction (e.g. pile driving (JNCC 2010b)) and decommissioning (e.g. use of explosives (2010c)). As part of the licencing procedures, developers and operators are required to utilise JNCC guidelines to minimise the risk of injury to cetaceans when undertaking such activities (JNCC, 2010b, 2010c; 2017; 2023; 2025; JNCC, Natural England & Cefas, 2025). The guidelines advise on conducting marine mammal observations prior to and during the activity and, where suitable, utilising procedures such as soft start (gradual introduction of the sound) to reduce and avoid direct harm to animals. A review of the marine mammal observer data demonstrated the effectiveness of soft start approach (Stone et al, 2017).

9.5: List of main conservation measures

MC03 Adapt/manage renewable energy installation, facilities and operation (excl. hydropower and abstraction activities). Guidance for the protection of marine European Protected Species from deliberate injury, killing and disturbance has been drafted (JNCC 2010a; Marine Scotland, 2014). Marine Industries generate a variety of noise through activities such as geophysical surveys (e.g. seismic surveys (JNCC 2017)), construction (e.g. pile driving (JNCC 2010b)) and decommissioning (e.g. use of explosives (2010c)). As part of the licencing procedures, developers and operators are required to utilise JNCC guidelines to minimise the risk of injury to cetaceans when undertaking such activities (JNCC, 2010b, 2010c; 2017; 2023; 2025; JNCC, Natural England & Cefas, 2025). The guidelines advise on conducting marine mammal observations prior to and during the activity and, where suitable, utilising procedures such as soft start (gradual introduction of the sound) to reduce and avoid direct harm to animals. A review of the marine mammal observer data demonstrated the effectiveness of soft start approach (Stone et al., 2017).

9.5: List of main
conservation measures

MG01 Management of professional/commercial fishing, shellfish and seaweed harvesting (incl. restoration of habitats). Fisheries Management Plans (FMPs) are currently being developed across all administrations for fisheries with perceived threats or pressures to the marine environment. FMPs are required under the Fisheries Act 2020 which provides the framework for management fisheries outside the EU Common Fisheries Policy. The Joint Fisheries Statement (agreeing the delivery of the 8 objectives of the Fisheries Act 2020) sets out plans for 43 FMPs. Publication of FMPs started last year and is expected to continue for 2-3 years. Some are being jointly developed, others by a single authority for its own waters. 6 FMPs have now been published.